Q1 - Define Tillage, objective & Type & Factor affecting of tillage

Def Tillage > Tillage is the manipulation of soil with tools and implements for loose the surface and bringing about conditions favourable for germination of seed & growth of crop.

Mechanical manipulation of the soil for preparing soil for growing & raising crops.

* Objective of Tillage >

- (1) To make the soil loose and porous
- (2) To derate the soil
- (3) To have repeated exchange of atmospheric air with the soil air
- (4) To increase the soil temprature
- (5) To control weeds
- (6) To remove stubble of the previous crop.
- (7) To destroy Insects.
- (8) to break hard pan.
- (9) To incorporate organic manufes and feetilizers into the soil
- (10) To invert the soil to improve Fertility (11) To prepare seed beel For germination of seed

TAGROIII

* TYPES OF TILLAGE >

from start of crop season to
the crop harvest known as "on
season tillage operation".

A preparatory fillage ->

- 1) primary tillage
- 11) secondary tillage
- 111) Layout of Seedbed & sowing
- B] seed bed preparation -
- or after tillage -

[II] off season tillage >

rillage operation during uncropped season for special purpose other than that for immediately raising the crop in the season are said to off seso season.

A preparatory tillage >

Tillage operation, which are carried out before the land is made ready for sowing or before sowing of crop is called "preparatory Tillage"

the following are preparatory Tillage.

- 1 Ploughing _
- 3 Manure mixing -
- @ clod crushing.
- 6 Compacting the soil-
- 3 leveling of land -
- (4) Harrowing -

[B] Seed bed preparation -

After preparatory tillage the land is to be laid out properly intigation of crops, if water, available for intigation and for sowing seeds, planting transplanting seedling. This operation is known as "seed bed preparation"

Thus Pollow of seed bed consist of -

- 1) Harrowing -
- 11) preparation of irrigation layouts -
- III) sowing -
- IV) covering seeds -

Intertillage or interculture.

- The tillage operations which are carried out in standing crop or in between crop rows are called as "inter culture"
- Inter tillage includes the operation -
 - (1) Gap Filling -
 - (11) Thinning -
 - (11) Hoeing -.
 - (14) Weeding -
 - (V) Top dressing of fertilizers -
 - (VI) Earthing up -
 - (VID Mulching -

Agronomy or Pholain it's Relationship other science

> Per Agronomy

Agronomy is the branch of Agriculture science which deals with the principle & practices of crop production & field management.

It is the study of plants in relation to soil and climate.

* Scope of Agronomy >

- (1) Agronomy is a dynamic discipline and scope of Agronomy is very vast
- (2) Agronomy is also concerned with the management of livestock including their feeding.
- (3) Agronomy also involves agronomic research on crops under different environmental condition like varying soil, climate, irrigation, feetilizer etc.
- (4) Agronomy also involves conducting the experiments in the field crops and laboratories.
- (3) It is also involves the application of research outcome into develop suitable package of practices for particular crops under given set of soil and climatic condition
- (1) It includes art of soil management.

 crop production, proper method of
 Tillage suitable period of cultivation.

* Relationship of Agronomy with other sciences

I] Basic science - It is science, which reveals
the facts or secrets of nature
e.g chemistry, physics, mathematics
Botany, Zoology

Applied Science - In the science in which
the theories and theories
and Laws propounded in
basic science are applied
for problems in agriculture
4 other fields e.g Agronomy,
Agricultural chemistry, Agricultural Entomology, AgricuItural Botany.

11 Agricultural chemistry ->

- comprising soil, water, plant Fertilizer and agril chemistry has been developed from basic science of chemistry

12 Plant pathology & Entomology >

- Development from botany and zoology, plant pathology is related with Agronomy for management of diseases.

3 Agricultural Economics

- Development from economics and useful for maintaining farm awards farm account and marketing

[4] Agricultural Extension >

-Development from sociology psychology and anthopology and anthopology and mainly related with different methods.

13 Agricultural Botany ->

- pevelopment from Botany and zoology and includes plant morphology plant physiology and plant breeding

16 Agricultural Engineering

- It is the concerned with care and use of improved tools implements and farm mechanical Machianary required for carrying out various field operation
- 1931 Define Weed & characteristics of Weed, and method of Weed control
 - > Def Weed >

Weed is a plant growing at a place and time where it is not desired

A weed is a plant growing where it is not wanted,

* characteristics of weed>

- (1) The weed seeds germinate early and the seedlings grow faster
- (2) They being hardy complete with crop plants for plant Nutrients maisture, space of Lower crop yields.

- (3) The flower earlier runs to seed in profusion and meture ahead of Therefore weeds control.
- (4) many weeds are non useful unwanted and undisirable.
- (3) Weeds are harmful to crops same of the weeds are also Harmful to cattle on human beings
- (6) They can thrive even under adverse conditions of soil. Climate Biotics stresses.

* Method of Weed control >

- (1) preventixe measures ->
- (11) control measures ->
 - a) Mechanical Methods >
 - b) cropping or cultural methods >
 - of Biological methods ->
 - d) chemical methods >
- (111) Integrated weed Management >

(1) preventive Measures >

- (1) use clean seed free from weed seeds
- (2) Use well decomposed compost or F.Y.M.
- (3) Destroy weeds before Flowering and seedling
- (4) Remove weed growth
- (5) Do not permit the livestock to move from weed infested area to clean area
- (6) Avoid shifting of soil from inferted

of seed.

Dep seed >

Any material use for planting and propagation, whether it is an the form of seed of food, fooder. Fibre and vegetable crops or seedling

OR

The seed may be defined as a feetilized oxule consisting of intact, embryo, stored food and seed coat which is viable & has good a capacity to germinate

* characteristics of good quality of seed.

- (1) It should be genetically pure
- (2) should exhibits or bear the morphological characters of the particular variety or hybrid of the crop
- (3) Have High germination percentage.
- (4) It should be dry and not mouldy in case of cereal, pulses, oil seed crop etc.
- (3) Give strong and vigorous seedling growth under normal condition
- (6) Well developed, plumpy, bold, uniform in shape, size, colour and texture.
- (7) clean, free from any admixture, dirt and inert makerial.
- (8) Free from naxious or objectionable or satellite weed seeds.
- (9) Dry and without moulds
 (10) Adapteeble crop variety or hybrid fitting
 in the cropping system.

What do you mean by cropping system 9 state Types, classification of cropping system - cropping system cropping system may be defined as the order In which the crop are cultivated on a piece of land over fixed period. * Types / classification of cropping system > I) Monoculture -In following or follow in rotation -III) Multiple cropping a) parallel multiple ctopping b) sequential multiple cropping a] parallel Multiple cropping > i) Mixed cropping ii) Intercropping iii) Relay cropping iv) Allay cropping V) multi stored cropping 5] Sequential Multiple cropping) i) <u>Sequence</u> cropping 9) Double b) Triple c) Quandicate ii) Ratoon cropping or Ratooning cropping system based on the following factors (1) This interest flight in the land (2) Extend of Land. (size of holding).

- (3) Nature of different holding Soil & their extends.
- (4) climatic co-ordination -
- (3) Availability of irrigation facility -
- (6) Agronomic character of crop.

properties of soil

> per Joil-

soil refers to the loose surface material of earth surface derived from the original rocks and minerals through weathering process,

* Physical properties of soil

- 1) soil texture -
 - 2) soil structure -
 - 3) porosity -
 - 4) BWK density -
 - 5) soil water -
 - 6) soil Temprature -

(1) Soil texture

- Texture refers to the composition with respect to relative proportion of sand, silt and clay in the soil
- sand partical ranges from 0.02 mm in diameter, silt particles from 0.02 mm to 0.002 mm and clay particles are less than 0.002 mm.
- If the soil contains more than 80% of silt, the soils are coursed as silty.

(2) | Soil structure |

- The arrangement of individual soil particles with respect to each other into a pattern is called soil structure.
- soil structure influences soil properties such as soil Erodibility, porosity, hydrawic conductivity, infiltration, erosion, capillary conductivity

(3) Porosity

- It is defined as the ratio of the volume of pores to the total soil volume.
 - Pores are macro (more than 0.6 mm diameter noncoupillary) and micro (less than 0.6 mm diameter coupillary)
 - sandy soil have 30% to 40% and clay soil have 50% to 60% pore spaces.
 - Addition of organic matter in the soil increases the porosity.

(4) Bulk density

- (gram/cm3) It is mass of oven dry soil per unit volume of soil.
- It is a apparent density.

(5) Soil Temproduret

- Tillage is a helpfel from maintaining optimum soil temprature.
- Tillage is useful For prepair air of water Relationship in the soil
- exparing the soil to hear of the sun.

- (6) (5011 water) ->
 - Tillage is useful for increasing Pnfiltrates moisture retentive capacity of the soil due to increase in pore spaces.
 - It also enable the free drainage upto water tube.

Define seed dormancy, white a Notes on seed dormancy & white it's types.

-> Def seed dormancy >>

seeds can be dormant if they are viable, but do not germinate even under favourable conditions

- (i) seed dormancy is helpful because it prevents pre-harvesting sprowting.
- (ii) If the crop is cought in roins at maturity stage, the seed will not germinate.
- (III) case of no dormancy the seeds will germinate in the field if cought in rains maturity
- (IV) It is also a bad character as seeds can not be used for so using immediately unless this period is over by breaking dormancy.

* Types of seed dormancy;

- (1) Innate dormancy ->
- (2) Enforced dormancy ->
- (3) Induced dormancy ->

II Innute Dormancy >

- May be due to the genetical characters of the seed or due to hard seed coat.

[2] Enforced dormancy >

- Due to the conditions of deficient oxygen, excess co2, deep placement.

[3] Induced dormancy -

- This Type of dormany results due to sudder physiological change in seed.

Q8] Define seed vaibility & explain it's Types or write down note on "seed vaibility"

> Def seed valability >

seed vaibility means that a seed is capable of germinating and producing a notmal seedling.

It is the ability or capacity of the seed to germinate.

* Types of seed vaibility >/ stages

- (1) Nucleus seed -
- (2) Breeder's seed -
- (3) Foundation seed -
- (4) Registered seed -
- (5) certified seed -

(1) Nucleus seed 7

- i) It is produced and maintained by the respective plant breeder at the main research station.
- Individual plants with true morphological and genetical character of variety of perticular crop.
 - tically purity and being very small in quantity are often costly.

(2) Breeder's seed

- i) It is very important class of seed ii) They are the seeds or vegetative propagation materials directly produced or controlled by originating plant breeder.
 - research steetion and if required on other research farms, agricultural college farms etc.

(3) Foundation seed

- i) The seed certification authority staff insect the plot from time to time.
- in the off type plants, disease and pest affected plants and weeds are removed for avoiding their mixture in seed.
- 111) As per seed act, the seed is tested for it's purity, germination and moisture percentage

(4) Registered seeds

- Registered seeds are the progeny of the foundation seeds, which are multiplied on the farms of registered under guidance and supervision of the seed certification staff.
- -IF the quantity of foundation seed is less than requirement then this stage of seed multiplication is followed.

(5) (Certified seeds)

- The Term certified seed production is widely used to denote the production of commercial seed solds to the formers for raising crops.
- The seed is tested for its purity germination and moisture content as per the seed certification / standards.
- Define Harvesting & Explain the method of Harvesting.
 - > Def Harvesting >

The process of separating crop plants from the Field is known as "Harvesting.

* Method of Harvesting >

· 1] cutting the plants close to the ground level >

In this method the cereal erops like Jawar, Bajara, wheat maize and paddy are cut close to the ground level by sharp sickles

(2) picking of pods or Fruits ->

- pods of the pulses crops
like mug, udid, pea, etc. are
picked up When they are
matured the fruits of Brinjal
tomato, bhendi and chilies
are also harvested by picking
method.

(3) pigging the produce from the soil >

- crops like turmeric, ginger, groundnut, sweet potato, onion and garlie are harvested by digging the soil with the help of kudali and produce is collected from the soil cleaned & stored.

(4) Mechanical Harvesting ->

-The groundnut and the potatto digger are used for harvesting of ground-nut and potato crop. By using the power machinery the crops like wheat and sugarcane also Harvested.

Q10] Write a Notes on "seed treatment"

-> * OBJECTIVE OF SEED TREATMENT ->.

- (1) control of Disease -
- (2) convenience in sowing -
- (3) <u>Quicker</u> germination —
- (4) Nitrogen fixation.

- (5) protection against insects_
- (6) Including earlines -
- (7) Inducing valiation -
- (8) Increasing yield -

* <u>Pifferent Field crops</u>

- 1) cotton cotton see are subjected to different treatment for easy and convenient sowing
 - ay cow dung paste treatment -
 - b) suphueic acid treatment -
 - c) cotton seeds are also treated with 1.1. mercurial compound such as agrosan @ 29 m/kg of seed for control of seed borne disease like wilt.
 - 11) cotiender > see are spited in two locules under slight pressure for even germination.
 - 111) Gaelic clove are separated before sowing in order to provide space
- QIII Give the classification of crops with suitable Example
 - => * classification of crops -
 - (1) classification on the basis of climate.
 - a) Tropical crop or warm season crops—

 The crops, which require

 warm climate or High temp

 are called Tropical crop

 e.g Rice, maize, sorghum.

- b) Temprate crops or cool season crops—)

 The crops which require cool and dry climate or low temp.

 and generally grown in winter season, are called temprate crops.

 e.g. Wheat, barley, oat, mastard.
- (2) classification on the basis of season y
 - i) kharif crop e.g tice, Ragi, maize, sorghum
 - ii) Rabi crop e.g wheat, barley, mustard
 - iii) <u>Summer crop</u> e.g summer groundnut, green gram, gourds, okrq etc
- (3) classification on the basis of life of a crop
 - i) <u>seasonal crop</u> e.g Rice, pearl millet, maize, sorghum, wheat, oat.
 - ii) Two seasonal crop-eig turmeric, ginger, long staple cotton.
 - iii) Annual crop e.g sugarcane
 - iv) Biennial crop- e.g Banand, papaya etc
 - v) perenial crops e.g. mango, guava, orange
- (4) classification on the basis of source of water
 - i) Rainfed crop -
 - The crop which is grow only on rain water is also Rainfed crop e.g rice, sorghum, maixe, pearl millete.
- (3) classification on basis of root system
 - i) Tap rooted crop-eig cotton, legumes, Red gram, black gram, cow pea
 - ii) Adventitious Root crop-e-g cereals, wheat, sorghum, maize

- (6) classification on the basis of economic importances
 - i) <u>- Cash crop</u> e.g coHon, sugarcane, coffee
 - ii) food crop-e-g Lucern, berseem, maize
 - (7) <u>classification</u> on the basis of use or Agronomic classification ->
 - i) <u>cereal or Grain crop-e-g</u> tice, wheat, maize, out battery
 - ii) <u>Legume or pulse crop-</u> e.g pigeon peg, black gram, horse gram
 - iii) forage or fodder crop-eig sorghum, maize, pajaro, cowpea
 - iv) Root crops e.g Radish, carrot, sugarbeet
 - V) Tuber crops e.g potedo, sweet potedo
 - vi) Fibre crop- e.g cotton, sunhemp, linseed
 - vii) <u>sugar crops-eig</u> sugar cane
 - Viii) oil seed or oil crops-eig castor, niger.
 - ix) Drug crops-eig tobacco
 - x) spice crops or spices > e.g termeric, ginger
 - xi) <u>Vegetable crops</u> reg Brinjal, tomato, cabbage, bhendi
 - XII) Green manute crops-e.g dhaincha

(8) Bolanical or Morphological or Taxonomic classification

- 1) Graminge-e.y All cereals, rice, maize, wheat
- ij Leguminoceae e.g All legumes, red gram
- iii) <u>cruciferaceae</u> e.g mustard; radish
- iv) <u>cycurbiticede</u> eg: cycumber, pumpkin
- y) Liliacege -eg. opion, gartle.
- vi) solanacege e. q poterto, tobaco.
- vi) malavacege e.g. cotton, olera
- Vii) Compositap- sunflower, sufflower
- VIII) convolvulaceae > sweet potato

Explain the Method of sowing (1) Brod casting -(2) Prilling or line sowing_ (3) Pibbling (4) Transplanting (5) planting (6) pulling seeds in plough furrow > (1) Brodusting It broad casting methods, seeds are spread uniformly over well-prepared land and is cover by ploughing or planking It is scattering or spreading of seed on the soil by band all over the field of then covering by soil with light Implements. tine sowing ? prilling or In this method, the seeds are (3) Dibbling placed in a furrow, pit or hole at a predermined spacing with a dibbler, planter or more commonly by hand . This method is ideal for crops such as groundnut, castor, maire, sugarcane, potato, onton - This method is suitable for crops requiring specific geometric area for canopy development. PHIlling or line sowing - To overcome the problems of broad asting drilling the seeds in line come into practice

- coxeting of seeds is necessary when indigeneous or two-bowl seed drill is used
- cost of mechanical seed drill is more
- This adopted for sowing crops like sorghum, pearl millet, upland rice, wheat, out.

(4) Transplanting)

- When more than one crop is to be grown in a year on the same piece of land, the time occupied by each crop has to be reduced.
- The seedling growth in the early stages is very slow.

(5) Planting

- some of the crops ar sown by using vegetative plant parts and the method of sowing in such crop is known as planting.
- preparation of inigation layouts is essential for planting & giving imaget to the crop.

(6) Putting seed in Plough Furrow

- This method is followed for erops like chickped and wal in some areas for better willization of residual soil moisture.
- The seeds are dropped behind the plough in the plough furrow with the belp of manual Labour and are covered by Successive turn of ploughing

*@13]. Ag1	roclimatic zones of Ma	hardshtra.
→ * A	GROCLIMATIC ZONES OF MAI	HARASHTRA >
(2) NOTH	- h konkan Costal zone - h konkan costal zone -	
(4) West (5) Sub (6) Wes (7) Cent	tern Ghot zone tern Maharashtra plain zon montane zone tern Maharashtra Scarcity = ral Maharashtra plateau zo	- Kolhapyr cone - Solapyr
	tral vidarbha zone —	Ya watmal sindewai .
>* [Ro] (D) A	of Agromist in Agriculties e of Agronomist > person who expert in A derivation of Agronomist'	gronomy is known
	the role of Agronomist in roportant	Agriculture is very

(3) Agronomist aims at obtaining maximum production at minimum cost.

(4) knowledge used for higher ctop production

- (5) He is concerned with production of food, fibre to meet the needs of increasing population.
- (6) Agronomist carry out research by considering soil, climate, crop/variety

(7) He is key person with working knowled de of all agricultural diciplines.

Enlist different tillage implements & explain the Factor cuffecting tillage operation.

Tillage implements >

(1) Wooden (indigenous or deshi) plough
(2) Mould board plough
(3) Disc plough puddler
(4) Chisel plough puddler -

(2) Mould board plough —

(3) Disc plough —

(4) Chisel plough puddler —

(5) Rotary plough or Rotary hee —

(6) Cultivators Blade harrow —

(7) puddler —

(8) Iron or inversion plough —

(9) Subsoil plough —

(9) subsoil plough (10) Ridge Plough (11) Blade harrow -

(12) Norwegian harrow -

* Factors offecting the tillage,

(1) Previous crops

- The types of crop decides the type and extent of preparatory cultivation required by the land.

- If the land is follow for several years and to bring under ewithout on deep tillage with is required.

(2) Type of soil =

- Fine textured soil like clay, poorly drained soils need deep cultivation to improve circulion, course sandy or loamy soils may not need deep cultivation

- Weed infested soil needs deep cultivation

(3) Weed Intensity

- Weed in Fester Fields needs deep (cultivation similarly land infuted perenial deep rooted weeds like kans etc.

- (4) [crop to be grown ->
 - peep cultivation also depends upon types of crop to be grown.
 - The crops which requires firm seed bed and which fibrous roof system like jawar, do not deep cultivation

(3) climate)

- climate inflyence the moisture content in the soil.
 - In low rainfall area deep cultivation is necessary to conserve more moisture

(1) Types of farming >

- crop under rainfed area are fully dependent on rain water.
 - only one crop during a year is grown.
- affecting soil productivity and Enlist the factors
 - > Def [soil productivity]

It is the capacity of the soil to produce crop with specific systems of management and its Expressed in terms of rields.

* Factor affecting soil productivity >

A Internal factors -

- the genetic or hereditary factors which can not be Manipulate.

0] External factors -

(1) precipitation -	N.A.
(11) Temprature -	
(11) solar radiation -	
(In Wind velocity -	
(10) Inind velocity - (v) atmospheric gases -	
(2) <u>Eduphic factors</u> —	
(3) BioHc factors -	
1) Plants -	
11) Bacteria -	<i>↓</i>
111) Animals -	
(4) physiographic factors -	
(3) Anthropic factors -	
Q17/ Define feetilizer. Enlist différent	Methods of
Fertilizer application.	
-> Deforethizer ->	
Any natural or mo	anu factiled
material dry or liqu	
the soil in order t	
or more plant nutti	ente
	O ID.
* Method of Feetilizer application,)
(1) Solid forms -	
A] Broadcasting - At sowin	nd and top draing
placement method - plous	7
8 wb so	,
of Localized Placement -	11, ~ ~ ~
- Contact	
- Pellet	
- Side diresing	
-band - Hill a	ind rows,

B] Liquid forms ->

- starter solution - folior application - direct application to soil - application of Feetilizer through

Integration water.

218] Difference between soil feetility 4 soil productivity
4 Intercropping & mixed cropping

> 1 [soil feetility]

- i) It is considered as an index of available nutrients to plants.
- If is one of the Fuctor for crop production
- III) It can be analyzed in the laboratory
- of the soll to produce crops

[Soil productivity]

- 1) It is a broader term to indicate yield of crops
- 11) It is interaction of all factors that determine the magnitude of yields
- Field under particular dimetic condition
- M It is resultant of maxing factors influering soil management.

1 Intercropping

- I) The main object to willize the space left between two rows of main crop.
- I) There is no competition between main 4 sub crop
- 111) subsiderly crop are of short duration and they are harvested eather than main crop.
- in rows and sowing

Mixed cropping

- 1) The main object is to get at least one crop under any climatic hazard.
- 11) Almost all crops compete with one another
- in The crops are almost same duration
 - In the crops may be broad castel.

@197 Give deteil classification of manuay and feetilizers with suitable example

) Bulky organic Manures

Bulky organic manures contain small percentage of major (N,P.+K) Hutelent they are applied in large quantity

2) Concentrated organic Manury

Concentrated organic manures Have higher percentage of major nutrient content than bulky organic manure.

- 1) <u>plant origin</u> d) Non edible oil cake b) Edible oil cakes
- 11) <u>Animal origin</u> e.g Blood meal, Aish meal

* Feetilizen

feetilizers are industrially manufactured chemicals containing plant Nutrients. Nutrients content is higher in feetilizer than in organic manures and nutrient availability is faster.

fclassification of feetilizers

- (1) straight feetilizers > e.g. Urea, Ammonium sulphate
- (2) complex feetilizers, contain two or three primary plant Nutrient. The primary Nutrients are chemically combined eg Diammonium phosphate, nittop-phosphate,
- (3) Mixed feetilizers > are physical mixture of straight feetilizers. Feetilizers are also grouped on the Nutrient in the Feetilizers.

a) Nitrogenous feutilizer

-1) <u>Nitrogen fettilizers</u> -> e.f. sodium nitrate, roleium Nitrate,

- 11) Ammoniacal feetilisers eng Ammonium sulphate, Ammonium chloride.
- 111) Nitrate Ammoniacal feetilizers e.g. Ammonium nitrate, calcium amonium nitrate

 (N) Amide feetilizers e.g. urea, calcium aynide.

DIPhosphatic feetilizers]

- 1) phosphatic feetilizer containing water soluble.

 phosphatic acid e.g. single superphosphate,

 double super phosphate
- 11) phosphalic feetilizer containing citéc acid soluble phospholic acid e.g basic elag.

111) phosphatic Feetilizer Containing Phosphoric acid.

@ Potash feetili 2015)

- potash or potassium containing feetilizer is called potassic feetilizer.
- chorde form e.g potassium chloride, mop
- 1) Non chiloride form e-g potassium Nitrate

Q20] perine Tilth, give the characteristics of 900d Tilth.

per Trilth !

It is the physical condition of soil resulting from Allage.

- * characteristics of good tilth }
- soil should be loose, porous, friable and free drainage.

- Micro and macro pores should be in equal proportions which facilities sufficient comount of moisture
 - size of soil aggregates is about 1-6 mm
 - It should be favourable for proper germination of seeds, emergence and proper growth and development of crop:

Q21] Difine Tillage. Effect of Tillage

per Tillage > Tillage is a manipulation of soil with tools and implements for loose the surface and bringing about conditions favourable for germination of seeds.

* [Effect of Tillage >[A] Beneficial Effect

(1) Effect on physical properties of the soil
a) pore space
b) soil structure
c) Bulk pensity
d) 90il water

el soil temprature -

(2) Effect on ehemical properties.

(#1) It a crelerates weathering of soil—

(#1) Leaching of toxic soils—

(#1) Improve the availability of Nutrients—

(IV) Increase efficiency of applied material—

(3) Effect on Biological properties—

1) Management of weeds

11) Management of insect pest of Pisease.

111) Proper functioning of plant roots.

Effect of Tillage on crop growth -(4) 1) Improve aeration ii) make the soil loose & poroug with increase. in Altration 111) Destroy insect / perts IN) Improve Structure of soil V) making favourable conditions for crop growth [2] Bad Effects L 1) compaction of soil due to heavy machinery 11) excessive tillage formation of powdamy condition of soil Intertillage most damage crop. IV) Import tillage may effect on

- 0X - - 8X

crop yield

This Note Downloaded from WEBSITE

Visit For Other B.Sc AGRICULTURE Notes, Practical Manual, Question

Paper, Model Answer Paper, And other Agriculture Information

WWW.BSCAGRISTUDY.ONLINE